

PHCOG MAG.: Research Article

Potential Pitfalls on the Physiological (Osmotic Fragility) Properties of the Red Blood Cell: Action of a Homeopathic Medicine

Juliana Rebello Gomes de Souza Fontes, Livia Pinella Guedes Pereira, Rosana Batista da Costa Gomes, Lourival Dias Pereira, and Sebastião David Santos-Filho

Centro Universitário de Volta Redonda – UniFOA, Av Paulo Erlei Alves Abrantes, 1325 27251-970, Três Poços, Volta Redonda, RJ, Brasil.

e-mail: jurebello@yahoo.com.br; dra.rosanacosta@ig.com.br ; Fone: 55-021-24-8122-3275

ABSTRACT - *Hypericum perforatum* (*Hypericum* p.) is a medicinal herb worldwide, primarily as a containing many polyphenolic compounds, popularly called St. John's wort or in homeopathic nomenclature of thousand needles plant. Its active principle is the hypericin. *Hypericum* p. has been used in popular medicine since ancient times for several disorders such as skin wounds, eczema, burns, and diseases of the alimentary tract, insomnia, and mental illness, among others as well as in intolerable pains. The capability of red blood cells (RBC) to resist hemolysis characterizes what is called the osmotic fragility (OF) of the membrane. We have investigated the possibility that *hypericum perforatum* dinamized homeopathically is capable of altering the physiological properties as RBC-OF. Briefly, blood samples were incubated with a *hypericum perforatum* 6CH or with sodium chloride solution (0,90% NaCl) as a control for 60 minutes at room temperature. Samples (25µl), were gently mixed with hypotonic NaCl (from 0.12 to 0.90%), centrifuged and the supernatants were isolated to determine the optical density (OD) in a spectrophotometer (540nm). Statistical analysis was performed. Osmotic fragility was significantly ($p<0.05$) changed by the presence of the *Hypericum perforatum* 6CH in the isotonic concentrations. Finally, the alterations to RBC membrane transport mechanisms were sufficient to promote modifications in osmotic fragility. Our results suggest that, the chemical results of *Hypericum perforatum* 6CH could be responsible for the observed effects.

KEYWORDS - Osmotic fragility; Red blood cell; *Hypericum perforatum* 6CH; Homeopathic medicine.

INTRODUCTION

The use of natural products has increased in all over the world, and popular interest in complementary and alternative medicine (CAM) in Brazil is as large as it is in other countries (1-4) especially with regard to phytotherapy, homeopathy and acupuncture. Medicinal plants are used for the human being however several biological effects and the consequences for the health have not been well established yet. Many plants contain active substances that can induce biological effects and their frequent use has been correlated with a high incidence of diseases or undesired biological effect in the population (5-7). However, various active compounds derived from medicinal plants have been assessed for their efficacy and tolerability in the treatment of breast cancer and have been evaluated in clinical trials (8).

Hypericum perforatum (*Hypericaceae*) is a medicinal plant species containing many polyphenolic

compounds, popularly called St. John's wort or in homeopathic nomenclature of thousand needles plant, it is a vivacious plant of the family of the *Hypericaceae*, who grows in hills and dry places, whose active principle is the hypericin. Has been used in popular medicine since ancient times for several disorders such as skin wounds, eczema, burns, and diseases of the alimentary tract, insomnia, and mental illness (9, 10), among others as well as in intolerable pains (10).

Hypericum p. extract contains flavonoids such as rutin, quercetin, and quercitrin, which demonstrated a free radical scavenging activity in a model of autooxidation of rat cerebral membranes (11). *Hypericum* p., in line with popular credence, also possesses anxiolytic, antiviral, wound healing, antimicrobial, analgesic, and anti-inflammatory effects (12).

Hypericum p. extract has always been referred to as having a benign side-effect profile compared to

tricyclic antidepressants and serotonin-specific reuptake inhibitors (13). Hypericum p. extract, as an efficacious antidepressant medication with a potential antioxidant activity, was therefore hypothesized to be useful in the treatment of pathological situations in which ROS play an important role such as acute inflammation (14).

Homeopathy, recognized as the medicine of the similarities, for using medicines that more resemble the patient; it was enunciated by Hipócrates (450AC): "Similia Similibus Curantur"; or either, *'Like Cures Like'* (15). Introduced by the German physician Samuel Christian Frederick Samuel Hahnemann at the beginning of the 19th century, consists of the treatment of the symptoms of a disease by means of ultra-diluted substances. These substances are chosen based on the assumption that they would cause the same symptoms in healthy subjects if ingested in measurable amounts. Minerals, vegetables, or substances obtained from living organisms may be used in the preparation of homeopathic medicines. Supposedly, the effect of these substances becomes stronger the more they are diluted. An important step in their preparation is called the *succussion*: "shakes" that are applied to the substance after each dilution stage. (16,17).

The homeopathic performance of Hypericum p. has been observed in the terminations of the sensitive nerves, in intolerable pains (10). Hypericum p. has been considered by some authors as antidepressant, so the homeopathic medicine is not used specifically for this. But considering its pathogenesis, described in the homeopathic materia medica, Hypericum p. has indicative to depression state (18).

Blood contains many types of cells with very different functions, ranging from the transport of oxygen to the production of antibodies. The sodium-potassium pump has a direct role in regulating red blood cell (RBC) volume: It controls the solute concentration inside the cell, thereby regulating the osmotic forces that can make a cell swell or shrink (19).

The capability of RBC to resist hemolysis characterizes what is called the osmotic fragility (OF) of the membrane. The osmotic fragility is classically used as a general screening procedure (20,21). The "fragility curve" reflects the structural and geometrical changes in RBC. Hemolytic results from a structural perturbation of the RBC and its cytoskeleton caused by its high partition in the membrane (20, 22).

The aim of this study was investigated the possibility that a medicine homeopathic Hypericum p. 6CH

(diluted and dinamized homeopathically) is capable of altering the physiological properties as well as osmotic fragility of the red blood cells (RBC).

MATERIAL AND METHODS

Reagents

The reagent NaCl (Merck S.A., Brazil) was used to prepare the solutions to evaluate osmotic fragility. Hypericum p. was obtained of the Novo extrato Farmácia homeopática Ltda, Brazil as a NaCl 0.90% dilution. Heparinized whole blood was withdrawn from a female sheep. The osmotic fragility evaluations of the RBC were performed with blood samples incubated with Hypericum p. dinamized homeopathically as 6CH.

The experimental procedure:

Blood was withdrawn from a female sheep with a heparinized syringe. The osmotic fragility evaluations of the RBC were performed with sheep blood samples incubated with Hypericum p. 6CH or with sodium chloride solution (0.90% NaCl) as a control for 60 minutes at room temperature. Briefly, RBC samples (25 µL), treated or not, were gently mixed with hypotonic NaCl (from 0.12 to 0.90%) solutions according to Dacie's modified method (Cavalcanti *et al*, 2003). After 30 min, at room temperature, these tubes were centrifuged (3500 rpm, 15 min). The supernatants were isolated to determine the hemoglobin optical density (OD) in a spectrophotometer (540 nm). The optical density of each supernatant was compared with that corresponding to stronger hypotonic solution (0.12% NaCl) that was considered 100% of hemolysis. The supernatant of the tube, which contained 0.90% NaCl was considered the "blank tube" for the reaction, because it has no hemolysis.

There are three intervals were determinates: interval 1 between 0.12 and 0.36% NaCl, interval 2 between 0.36 and 0.60% NaCl and interval 3 between 0.60 and 0.90% NaCl, according the curve tendency (Cavalcanti *et al*, 2003). The means and SD of each interval was determinate and the statistical analysis was performed. The experiments were carried out in compliance with guidelines on the use of live animals in scientific investigations.

Statistical analysis

The results were compared with the control samples, and statistical analysis was performed by independent test ($p < 0,05$) to determine the significance of the difference between incubated with Hypericum p. 6CH and control samples.


RESULTS

Figure 1 shows the osmotic fragility of the sheep erythrocyte incubated with Hypericum p. 6CH when

reacted with different NaCl hypotonic solutions. The curve tendency shows that in hypotonic and isotonic solutions the hemolysis increased significantly ($p < 0.05$). Figure 2 presents the mean of the osmotic fragility after analysis of the three NaCl concentrations intervals obtained of the osmotic curve of the figure 1.


The analysis of the results confirmed a significant statistical increase ($p < 0.05$) on osmotic fragility of the erythrocyte incubated with Hypericum p. 6CH in the intervals 2 (0.36 to 0.60% NaCl) and 3 (0.60 to 0.90% NaCl).

Figure 1: Osmotic fragility tendency of red blood cell, treated with *Hypericum perforatum*



The osmotic fragility evaluations of the RBC were performed with blood samples incubated with hypericum perforatum 6CH (diluted and dinamized homeopathically) or with sodium chloride solution (0.90% NaCl) as a control, for 60 minutes at 37°. Briefly, RBC samples (25µl), treated or not, were gently mixed with hypotonic NaCl (from 0.12 to 0.90%) solutions according to Dacie's modified method. After 30 min., at 37°, these tubes were centrifuged (1500rpm, 15 min). The supernatants were isolated to determine the optical density (OD) in a spectrophotometer (540nm). The optical density of each supernatant was compared with that corresponding to 100% lyses (0.12% NaCl). The supernatant of the tube, which contained 0.90% NaCl, was considered the "blank tube" for the reaction.

Figure 2: The osmotic fragility of red blood cells treated with *Hypericum perforatum*



Data in each interval represent mean $\pm$ SD. Three intervals were determinates: interval 1 between 0.12 and 0.36% NaCl, interval 2 between 0.36 and 0.60% NaCl and interval 3 between 0.60 and 0.90% NaCl, according the curve tendency. The means and SD of each interval was determinate and the t-test. ($p < 0.05$) was used to analyze potential differences between the percentage of hemolysis of each interval.

DISCUSSION

Many authors have cited a large number of drugs which cause alterations on the shape and physiology of the red cells (23, 24). The evidence that drugs can interfere on the labeling of RBC has been demonstrated with various natural products, like *Thuya occidentalis* (25), tobacco (26), *Peumus boldus* (27), *Maytenus ilicifolia*, (28), *Mentha crispa* (29), *Hypericum perforatum* (30).

Previously, we have reported that *Mentha crispa* and *Piper Methysticum* alter morphologically the RBC by expressing an increased morphometric factor (31). Moreover, the results obtained with the quality comparison of the shape of the RBC (non treated and treated with natural extracts) under optical microscopy could justify the modifications in the uptake of ^{99m}Tc by red blood cells in the presence of *Mentha crispa*

extract, similar to that observed with the extract of *Maytenus ilicifolia* (28). The achieved results have revealed important morphological alterations due to the treatment of blood with *Mentha crispa* extract (29). Either, *Hypericum p.* extract produced modifications in the uptake of ^{99m}Tc by RBC and in the uptake of the radiopharmaceutical sodium pertechnetate (^{99m}TcO₄Na) by thyroid, stomach and pancreas (30).

The homeopathic *Hypericum p.* has indicative to depression state (18), and intolerable pains (10). However, we could not find report of natural products diluted and dinamized homeopathically could interfere on the labeling of RBC.

Erythrocyte osmotic fragility is the resistance of RBC hemolysis to osmotic changes that is used to evaluate RBC friability (32). In the present study we have found that the osmotic fragility of RBC's was changed by presence of *Hypericum p.* 6CH in the studied concentrations. The results showed that there was a significant statistical ($p < 0.05$) increase in the osmotic fragility of those cells treated with *Hypericum p.* 6CH (71%) relative to control (68%) in the interval 2 (0.36 and 0.60% NaCl) that is hypotonic interval of the osmotic curve (Figure 1). In the interval 3 (0.60 and 0.90% NaCl) that is isotonic interval, the osmotic fragility also increased significantly ($p < 0.05$) with *Hypericum p.* 6CH (16%) relative to control (10%).

These experimental data show that the osmotic fragility of the RBC can be increased in the presence of *Hypericum p.* 6CH solution and we can suggest that this effect may be due to the properties of this medicine that may (i) alter the RBC physical properties, (ii) have

a direct or an indirect effect on intracellular sodium ion concentration and (iii) the medicine homeopathic could not be considered a placebo effect.

Conclusion

The experiment shows that the osmotic fragility of the RBC can be increased in the presence of the *Hypericum p.* 6CH and we can suggest that this effect may be due to the properties of this homeopathic medicine.

REFERENCES

1. P. Fisher and A. Ward. Complementary medicine in Europe. *BMJ*. **309(6947)**: 107-11 (1994).
2. D.M. Eisenberg, R.C. Kessler, C. Foster, F.E. Norlock, D.R. Calkins and T.L. Delbanco. Unconventional medicine in the United States. Prevalence, costs and patterns of use. *N Engl J Med*. **328(4)**: 246-52 (1993).
3. D.M. Eisenberg, R.B. Davis, S.L. Ettner, S. Appel, S. Wilkey, M. Van Rompay and R.C. Kessler. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA* **280(18)**: 1569-75 (1998).
4. M.A. Burg, R.L. Hatch and A.H. Neims. Lifetime use of alternative therapy: a study of Florida residents. *South Med J*. **91(12)**: 1126-31 (1998).
5. T. Nguyen, L. Fluss, R. Madej, C. Ginther and T. Leighton. The distribution of mutagenic activity in red, rose and white wines. *Mutat Res*. **223(2)**: 205-12 (1989).
6. J. Subiza, J.L. Subiza, P.M. Escribano, M. Hinojosa, R. Garcia, M. Jerez and E. Subiza. Occupational asthma caused by Brazil ginseng dust. *J Allergy Clin Immunol* **88(5)**: 731-6 (1991).
7. C.A. da Fonseca, J. Leal, S.S. Costa and A.C. Leitão. Genotoxic and mutagenic effect of guarana (*Paullinia cupana*) in prokaryotic organisms. *Mutat Res*. **321(3)**: 165-73 (1994).
8. D. Mantle, T.W. Lennard and A.T. Pickering. Therapeutic applications of medicinal plants in the treatment of breast cancer: a review of their pharmacology, efficacy and tolerability. *Adverse Drug React Toxicol Rev*. **19(3)**: 223-40 (2000).
9. J. Barnes, L.A. Anderson and J.D. Phillipson. St. John's wort (*Hypericum perforatum* L.): a review of its chemistry, pharmacology and clinical properties. *J Pharm Pharmacol*. **53(5)**: 583-600 (2001).
10. M.B. Argenta, *Matéria Médica Homeopática: sinais e sintomas odontológicos*. (Techmed, Riberão Preto, 2005) pp.____.
11. A. Saija, M. Scalese, M. Lanza, D. Marzullo, F. Bonina and F. Castelli. Flavonoids as antioxidant agents: importance of their interaction with

- biomembranes. *Free Radic Biol Med.* **19(4)**: 481-6 (1995).
12. O.A. Flausino Jr., H. Zangrossi Jr., J.V. Salgado and M.B. Viana. Effects of acute and chronic treatment with *Hypericum perforatum* L. (LI 160) on different anxiety-related responses in rats. *Pharmacol Biochem Behav.* **71(1-2)**: 251- 7 (2002).
13. B. Vitiello. *Hypericum perforatum* extracts as potential antidepressants. *J Pharm Pharmacol.* **51(5)**: 513-7 (1999).
14. M. Menegazzi, R. Di Paola, E. Mazzon, C. Muiá, T. Genovese, C. Crisafulli, H. Suzuki and S. Cuzzocrea. *Hypericum perforatum* attenuates the development of carrageenan-induced lung injury in mice. *Free Radic Biol Med.* **40(5)**: 740-53 (2006).
15. P.M. Zelante, R.O. Nardy, *Arnica e o processo inflamatório / Miasmas na cavidade bucal*, (LR, São Paulo, 2001) pp.
16. M.A. Sampaio. *Odontologia Holística - A Saúde Começa pela Boca*, (Mythos/IBHE, São Paulo, 1994) pp. .
17. R.M. Almeida. A critical review of the possible benefits associated with homeopathic medicine. *Rev Hosp Clin Fac Med Sao Paulo.* **58(6)**: 324-31 (2003).
18. F.C. Goulart. *Avaliação dos efeitos Hypericum perforatum dinamizado homeopaticamente em comportamento de ratos.* (VIII Simpósio Nacional de Pesquisa em Homeopatia SINAPIH e VIII Encontro Internacional de Pesquisa em Homeopatia, São Paulo, 2004).
19. B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, P. Walter, *Molecular Biology of The Cell*, (Garland Science, New York, 2002) pp 306-7.
20. J. Didelon, P. Mazon, S. Muller and J.F. Stoltz. Osmotic fragility of the erythrocyte membrane: characterization by modeling of the transmittance curve as a function of NaCl concentration. *Biorheology.* **37(5-6)**: 409-16 (2000).
21. X. Wang, L. Wei, J.P. Ouyang, S. Muller, M. Gentils, G. Cauchois and J.F. Stoltz. Effects of an angelica extract on human erythrocyte aggregation, deformation and osmotic fragility. *Clin Hemorheol Microcirc.* **24(3)**: 201-5 (2001).
22. T.C. Cavalcanti, C.C. Gregorini, F. Guimarães, O. Rettori and A.N. Vieira-Matos. Changes in red blood cell osmotic fragility induced by total plasma and plasma fractions obtained from rats bearing progressive and regressive variants of the Walker 256 tumor. *Braz J Med Biol Res.* **36(7)**: 887-95 (2003).
23. S. Ammus and A.A. Yunis. Drug-induced red cell dyscrasias. *Blood Rev.* **3(2)**: 71-82 (1989).
24. A.C. Braga, M.B.N Oliveira, G.D. Feliciano, I.W. Reininger, J.F. Oliveira, C.R. Silva and M. Bernardo-Filho. The effect of drugs on the labeling of blood elements with Technetium-99m. *Curr Pharm Des.* **6(11)**: 1179-91 (2000).
25. J.F. Oliveira, A.C. Braga, A.S. Ávila, L.M. Fonseca, B. Gutfilen and M. Bernardo-Filho. Effect of *Thuya occidentalis* on the labeling of red blood cell and plasma proteins with technetium-99m. *Yale J Biol Med* **69(6)**: 489-94 (1997).
26. M.V. Vidal, B. Gutfilen, L.M. Barbosa-da-Fonseca and M. Bernardo-Filho. Influence of tobacco on the labelling of red blood cells and plasma proteins with technetium-99m. *J Exp Clin Cancer Res.* **17(1)**: 41-6 (1998).
27. I.W. Reiniger, C.R. Silva, I. Felzenszwalb, J.C. Mattos, J.F. Oliveira, F.J.S. Dantas, A.A.C. Bezerra, A. Caldeira-de-Araujo and M. Bernardo-Filho. Boldine action against the stannous chloride effect. *J Ethnopharmacol.* **68(1-3)**: 345-8 (1999).
28. J.F. de Oliveira, A.C. Braga, M.B. de Oliveira, A.S. Ávila, A.C. de-Araújo, V.N. Cardoso, R.J. Bezerra and M. Bernardo-Filho. Assesment of the effect of *Maytenus ilicifolia* (Espinheira Santa) extract on the labeling of red blood cells and plasma proteins with technetium-99m. *J Ethnopharmacol.* **72(1-2)**: 179-84 (2000).
29. S.D. Santos-Filho, G.L. Diré, E. Lima, M.N. Oliveira and M. Bernardo-Filho. Effect of *Mentha crispa* (mint) extract on the labeling of blood elements with technetium-99m: A possible evaluation of free radicals. *J Biol Sci* **4(3)**: 266-70 (2004).
30. S.D. Santos-Filho and M. Bernardo-Filho. Effect of *Hypericum perforatum* extract on *in vitro* labelling of blood elements with technetium-99m and on biodisponibility of sodium pertechnetate in Wistar rats. *Acta Cir Bras* **20(Suppl 1)**: 76-80 (2005).
31. S.D. Santos-Filho, C.K. Ribeiro, G.F. Diré, E. Lima and M. Bernardo-Filho. *Technetium, Rhenium and other Metals in Chemistry an Nuclear Medicine* (eds) Nicolini M and Mazzi U (SGE editoriali, Padova, 2002) **6**:503-5.
32. S. G. Wu, F.R. Jeng, S.Y. Wei, C.Z. Su, T.C. Chung, W.J. Chang and H.W. Chang. Red blood cell osmotic fragility in chronically hemodialyzed patients. *Nephron.* **78(1)**: 28-32 (1998).