

**PHCOG MAG. SHORT
COMMUNICATION**
**Hepato-Protective activity of
Moringa oleifera Lam Leaves in
Carbon tetrachloride induced
Hepato-Toxicity in Albino Rats.**

D. Selvakumar ^{a,*}, P. Natarajan ^b

^aResearch and Development Wing, Department of
Quality control, Arya Vaidya Nilayam Limited,
Madurai., Tamilnadu, India.

^bDepartment of Pharmacy, Annamalai university,
Chidambaram, Tamilnadu, India.

*Correspondence : deselva123@rediffmail.com

ABSTRACT

The present study was conducted to evaluate the hepato-protective activity of methanolic and chloroform extract of *moringa oleifera* leaves are evaluated in CCl₄-induced hepatotoxicity in albino rats. Silymarin (200mg/kg) was given as reference standard. The methanolic and chloroform extracts of leaves of *moringa oleifera* have shown very significant hepatoprotection against CCl₄-induced hepatotoxicity in albino rats in reducing Serum total bilirubin, direct bilirubin, SGPT and SGOT levels.

Key words: *Moringa oleifera*, hepatotoxicity, Drumstick

INTRODUCTION

Liver disease remains one of the serious health problems. Herbs play a major role in the management of various liver disorders. A number of plants possess hepatoprotective property [1]. *Moringa oleifera* Lam (Moringaceae) commonly known as drumstick, is used in Indian folk medicine for the treatment of various illness. The leaves are rich in Vitamins A and C and considered useful in scurvy [2]. The flowers reported to contain quercetin and kempferol and used as uterus tonic [2]. The pods and leaves contain Vitamin A, B, E and C. They contain eight essential amino acids [2, 4,

5]. The flowers reported to contain quercetin and kempferol [2].

MATERIALS AND METHODS

Plant material

The plant material used in this study was collected during January 2005 in Chidambaram, a rural town of Tamilnadu, India and authenticated by herbarium staff of Department of Botany, Annamalai University in Chidambaram. A voucher specimen has been deposited at the herbarium of the department of pharmacognosy, Annamalai University in Chidambaram, Tamilnadu, India.

Preparation of the Extract

The shade dried leaves of *Moringa oleifera* Lam was extracted with chloroform and methanol successively by soxhlation method for 6 hours. Filtered, concentrated over water bath and evaporated under reduced pressure. The yields of methanolic extract (yield 7.4%) and chloroform extract (yield 8.6%) were calculated.

Animals

Albino rats (either sex) of Sprague dawley strain, weighing 150-200g were used. The animals were acclimatized to laboratory conditions (RT-25°C) for 4 days and given pelleted animal feed (Hindustan Lever) and drinking water, Diagnostic reagent kits (Enzopak) were used for the estimation of serum SGPT and SGOT levels.

Hepatoprotective Activity

The animals were divided into five groups comprising of six albino rats in each group using randomization technique and treated with the extract for seven days to assess the hepato-protective potential of the plant. The first group (vehicle control) received vehicle for all the seven days. The second group was kept as toxin control and given only the CCl₄ treatment. The third group received methanol extract in the dose of 500mg/kg p.o. and the fourth group received the chloroform extract in the dose of 500mg/kg p.o. The fifth group received Silymarin in the dose of 200mg/kg, p.o. as a reference material for the study. All the animals except the vehicle control received CCl₄ on 5th day of the treatment. The animals were sacrificed by cervical dislocation after 48 hours of CCl₄ administration. The blood samples were collected by cardiac puncture in heparinized microfuge tubes. The blood samples thus collected were immediately centrifuged at 2500rpm for 15 minutes. When serum clearly separated out, the serum is analyzed for SGPT and SGOT levels using enzopak reagent kits by the method proposed by Reitman and Frankel [10]. The

Table: 1 Effect of methanolic and chloroform extracts of moringa oleifera leaves on CCl₄-induced hepatotoxicity

Group	Total Bilirubin ^a (mg/dl)	Direct Bilirubin ^a (mg/dl)	SGPT (Units/ml) ^a	SGOT (Units/ml) ^a
Control	0.94±0.27	0.76±0.14	56.01±1.58	59.40±4.51
CCl ₄	7.05±0.66*	3.21±0.05*	114.40±2.03*	131.0±2.96*
Methanolic Extract (500mg/kg)	1.08±0.28*	1.17±0.06*	42.57±1.03*	51.0±0.68*
Chloroform Extract (500mg/kg)	1.22±0.27*	1.22±0.06*	54.51±1.59*	56±1.24*
Silymarin (200mg/kg)	1.22±0.13*	1.24±0.05*	38.76±1.32*	52.0±1.28*

^a Values of mean ± S.E.M. (n=6)

* P < 0.01 vs. control, Student's 't' test

results thus obtained were subjected to statistical analysis using student t-test and analysis of variance (Table: 1)

RESULTS AND DISCUSSION

The methanolic and chloroform extracts of leaves of *moringa oleifera* have shown very significant hepatoprotection against CCl₄-induced hepatotoxicity in albino rats in reducing Serum total bilirubin, direct bilirubin, SGPT and SGOT levels. Liver section of moringa oleifera treated animal group clearly showed normal hepatic cells and central vein thereby confirming hepatoprotective activity.

REFERENCES:

1. S.S. Handa, A. Sharma and K.K. Chakraborti. Natural products and plants as liver protecting drugs. *Fitoterapia*. 57: 307-45 (1986). CCl₄ induced hepatic injury in rat. *Ind J Pharmacol*. 33: 260-266(2001).
2. Anonymous, Raw Materials, the Wealth of India, Publication and Information Directorate (CSIR), New Delhi, 1985.
3. Hartwell, JL, Plants used against cancer, a survey, *Lloydia*, 30-34 (1967-1971).
4. Verma SC, Banerji R, Misra G, and Nigam SK. Nutritional value of *Moringa*. *Curr. Sci.* 45(21): 769-770 (1976).
5. Lowell JF, *Moringa oleifera*: Natural nutrition for the tropics, the miracle tree, Church world Service, Dakar, Senegal, 1989.
6. Ruckmani, K., Kavimani, S., and Jaykar, B., Effect of moringa oleifera lam on paracetamol-induced Hepatotoxicity. *Ind. J. Pharm. Sci.* 60(1):33-35(1998).
7. Kaushik, R., Rimi, H., and Debjani, G., Central inhibitory effect of *Moringa oleifera* root extract: Possible role of neurotransmitters. *Ind. J. Exp. Biol.*, 41: 1279-1284(2003).
8. Caceres, A., Saravia, A., Rizzo, S., Lorenza, Z., De leon, E. and Nave, F., Pharmacological properties of *Moringa oleifera*, 2: screening for antispasmodic, anti-inflammatory and diuretic activity, *J.Ethnopharmacol.* 36: 233-237(1992).
9. Rao, K.S., and Mishra, S.H., Anti-inflammatory and antihepatotoxic activities of the roots of moringa pterygosperma gaertn. *Ind. J. Pharm. Sci.*, 60(1): 12-16(1998).
10. Ashok, S.K., Somayaji, S.N., Bairy, K. L., Hepatoprotective effects of Ginkgo Biloba against CCl₄ induced hepatic injury in rat. *Ind. J. Pharmacol.*, 33, 260-266(2001).
11. Reitman, S., Frankel, S., A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Amer.J.Clin.Path.*, 28,56-63(1957).
