

## PHCOG MAG.: Research Article

# GC-MS and Bioactivity of the Essential Oil of *Senecio rowleyanus* Jacobs

El Hawary S. S., Galal A. E., Yousif M. F. and Kirolos F. N.

Department of Pharmacognosy, College of Pharmacy, Kasr El-Einy Street, 11562, Cairo University, Cairo, Egypt  
[miriamyousif@hotmail.com](mailto:miriamyousif@hotmail.com)

### ABSTRACT

The chemical composition of the essential oil of *Senecio rowleyanus* Jacobs obtained by hydro distillation was analyzed by GC-MS and tested for cytotoxic and antimicrobial activities. Twenty five components representing 99.95% of the oil composition were identified. The oil was found to contain a significant amount of sesquiterpene alcohols of the guaiane type. Spathulenol (22.9%), germacrene B (12.4%), myrcene (12.8%) and viridiflorol (11%), were the predominant components. The oil exhibited marked cytotoxic activity against certain brain and liver human cell lines *in vitro*. In addition, it showed noticeable antimicrobial effect against the test microorganisms.

**KEYWORDS:** *Senecio rowleyanus* Jacobs; Essential oil; Antimicrobial activity; Cytotoxic activity; Spathulenol

### INTRODUCTION

*Senecio rowleyanus* Jacobs (1) is an ornamental plant cultivated in several public gardens in Egypt. The essential oil of different *Senecio* species have been previously investigated (2-8). Several of these oils exhibited antimicrobial activity (6). It has been noticed that many recent publications focus on the antitumor activity of the essential oils of some plants (9, 10). A study of the alkaloid content and cytotoxic and antimicrobial activities of whole extracts of *S. rowleyanus* herb was presented in an earlier communication (11). However, reports neither on composition nor on the bioactivity of the essential oil of *S. rowleyanus* could be traced in the available literature. Thus, it was deemed of interest to pursue investigation of the chemical composition and the bioactivities of the essential oil of the title plant as described in this note.

### MATERIAL AND METHODS

#### Plant material

Samples of *S. rowleyanus* herb were obtained from plants cultivated in a private garden, Kaliobeya governorate, during the flowering stage, from February to May, 2004. Identity of the collected plant was kindly verified by Dr. M. El Guibaly, Researcher of Plant Taxonomy, National Research Center, Dokki, Guiza.

#### Preparation of oil

The fresh plant (500g) was subjected to hydro distillation for 6 hours, using a Clavenger-type apparatus. The oil was collected, dried over anhydrous sodium sulfate followed by GC-MS analysis.

#### GC-MS analysis

The oil was analyzed using a Hewlet Packard 5970 GC

coupled with a mass spectrometer equipped with FID and data system SS- 180. The capillary column (20 m x 0.2mm i.d.), coated with Carbowax 20 M and 0.33µm film thickness, was used. Carrier gas, helium; injection temperature, 200°C; oven temperature program; initial temperature was kept at 60°C for 3 min and programmed to 200°C at a rate of 3°C/min, and kept constant at 200°C for 15 min; flow rate, 1 ml/ min; final temperature, 200°C; detector temperature, 250°C; sample size, 0.5 µL of 10% solution in ether; separation oven temperature, 248°C; transfer line temperature, 260°C; analyzer pressure, 10<sup>-4</sup> torr; scan time, 2 sec/ scan; scan range, 20-550; E1+ QIMS LMR UP LR mode electron ionization voltage, 70e.v. and electron multiplier voltage 1800 e.v. Identification of the oil constituents was achieved by library search based on a Willey 275 L GC-MS library and by comparing their retention indices and mass fragmentation pattern to those of the available references (12-15). The quantitative estimation was determined by relative peak area measurement. A series of authentic *n*-alkanes was subjected to GLC under the same conditions; the retention time of each *n*-alkane was observed and Kovat's indices of oil constituents were calculated (15). Results are shown (Table 1). A fingerprint of the oil was obtained by plotting the Kovat's index of each oil constituent against its percentage in the oil (Fig.1).

#### Bioactivities of the essential oil

##### *In vitro* screening for cytotoxic activity.

Tumor cell lines: Liver carcinoma cell line (MCF8), and brain tumor cell line (U251) maintained in laboratory

of National Cancer Institute, Cairo, Egypt were used. The test solution was the previously prepared essential oil. Different concentrations in DMSO were used. The test solution was screened for cytotoxic activity adopting sulforhodamine B stain (SRB) assay (16, 17). Different concentrations of the tested solution (0-12.5 µg/ml) were added to the cell monolayer. Triplicate wells were prepared for each individual dose. Monolayer cells were incubated for 48 hours at 37°C and in 5% CO<sub>2</sub> atmosphere. After 48 hours, cells were fixed, washed and stained with sulforhodamine B stain. Excess stain was washed with acetic acid and attached stain was recovered with Tris EDTA buffer. Color intensity was measured by an ELISA reader. The relation between surviving fractions and extract concentration was plotted to get the survival curve of each tumor cell line after the specific extract concentration. The dose which reduces survival to 50% (IC<sub>50</sub>) and 10% (IC<sub>10</sub>) growth inhibition were calculated on the fore mentioned human cell lines. Results are recorded (Table 2).

#### Testing of the antimicrobial activity

The antimicrobial activity was carried out in Microbiology Department, Faculty of Pharmacy, Cairo University. Microorganisms used included: Gram positive bacteria (*Bacillus subtilis* (NCTC) and *Staphylococcus aureus* (ATCC)), Gram negative bacteria: (*Escherichia coli* (ATCC), *Pseudomonas aeruginosa* (CNCMA), *Proteus vulgaris* (NCTC), and *Klebsiella pneumonia*), acid fast bacteria (*Mycobacterium phlei*) and fungi (*Candida albicans* and *Saccharomyces cerevisiae*). Rifampicin and nystatin impregnated discs were used as reference standards.

The antimicrobial screening was performed according to Duguid *et al* (18) and Lorian (19). The oil was diluted and solubilized in dimethylsulphoxide (DMSO) at concentration series of 40%, 20%, 10%, and 5%. The antimicrobial activity was performed by disc agar diffusion method against reference strains of Gram positive bacteria, Gram negative bacteria and fungi. A disc impregnated with 20 µL of DMSO was used as a negative control. Rifampicin, and nystatin impregnated discs were used as positive control. Results are compiled (Table 3).

#### RESULTS AND DISCUSSION

The percentage of essential oil in *S. rowleyanus* was 0.1%. Regarding the physical characters, the distilled oil was a colorless liquid with yellow tint, pleasant aromatic odor, and acrid taste with refractive index 1.482.

GC/MS analysis of the essential oil of *S. rowleyanus* (Table1) revealed the presence of 25 identified components representing 99.95% of the oil composition. Study of the oil composition revealed the presence of a hydrocarbon fraction representing 54.44 % of the oil, besides an oxygenated fraction representing 45.51 %. The hydrocarbon fraction consisted of monoterpenes (26.44%), with myrcene (12.8 %) as a major component. Sesquiterpenes were also present at 28.02 %, of which germacrene B (12.4%) and *trans*-caryophyllene (8.42%) exist as predominant components. The oxygenated fraction comprised, predominantly, sesquiterpene alcohols (40.02%), with minor amounts of oxides (5.46 %). The alcohols detected in the oil were principally of the guaiane skeleton and included spathulenol (22.9%), as the major component of the oil followed by other alcohols, viridiflorol (10.99%), globulol (1.16 %), epiglobulol (0.86%), ledol (0.42 %), and carotol (1.16 %). Oxygenated monoterpene derivatives could not be detected.


Results revealed that the essential oil of *S. rowleyanus* showed significant cytotoxic activity against the growth of human cell lines (brain tumor cell line (U251) and liver carcinoma cell line (MCF8). It showed a marked activity against liver carcinoma cell line with IC<sub>10</sub> 9.5 µg/ml and IC<sub>50</sub> 5µg/ml. It even displayed a more pronounced activity against liver carcinoma cell line with IC<sub>10</sub> 8 µg/mL, and IC<sub>50</sub> 2.6 µg/mL. (Table 2). The observed cytotoxic activity of the essential oil may be attributed to the presence of some monoterpene hydrocarbons (α-pinene, β-pinene and limonene) which have been previously reported as cytotoxic components in a number of essential oils (20). Results are comparable to those produced by the essential oil of other *Senecio* species (21).

From table (3) it is obvious that the essential oil of *S. rowleyanus* inhibited the growth of almost all the test micro organisms in a dose dependant manner as compared to the control drug. The effect was more pronounced on Gram positive and acid fast bacteria than on Gram negative ones, however being only moderate against *Pseudomonas aeruginosa*. Results obtained correlate with those concerning the antimicrobial activity of other *Senecio* species (22). Moreover the marked activity on *Mycobacterium phlei* may suggest their use as for certain *Senecio* species in treatment of tuberculosis (23).

Table 1 : Results of GC-MS analysis of the essential oil of *S. rowleyanus*.

| Peak no. | $R_t^*$ | Mass spectral data |                   | Identified Component        | KI <sup>***</sup> | %    |
|----------|---------|--------------------|-------------------|-----------------------------|-------------------|------|
|          |         | M <sup>+</sup>     | B P <sup>**</sup> |                             |                   |      |
| 1        | 4.29    | 136                | 93                | $\alpha$ – Pinene           | 518               | 3.62 |
| 2        | 5.38    | 136                | 93                | $\beta$ – Pinene            | 626               | 3.13 |
| 3        | 5.58    | 136                | 93                | Sabinene                    | 647               | 1.51 |
| 4        | 6.33    | 136                | 41                | Myrcene                     | 720               | 12.8 |
| 5        | 7.06    | 136                | 67                | Limonene                    | 783               | 2.34 |
| 6        | 8.86    | 134                | 119               | <i>p</i> – Cymene           | 929               | 1.51 |
| 7        | 11.13   | 138                | 55                | <i>trans</i> -Pinane        | 1067              | 1.53 |
| 8        | 16.24   | 204                | 105               | $\alpha$ – Copaene          | 1272              | 1.65 |
| 9        | 20.33   | 204                | 41                | <i>trans</i> -Caryophyllene | 1381              | 8.42 |
| 10       | 22.02   | 204                | 41                | Isocaryophyllene            | 1449              | 0.63 |
| 11       | 22.99   | 204                | 93                | $\alpha$ – Humulene         | 1503              | 0.82 |
| 12       | 25.61   | 204                | 121               | Germacrene B                | 1641              | 12.4 |
| 13       | 26.33   | 204                | 41                | $\beta$ – Seline            | 1680              | 0.39 |
| 14       | 27.83   | 204                | 41                | $\beta$ – Bisabolene        | 1758              | 3.37 |
| 15       | 28.92   | 204                | 41                | $\gamma$ – Elemene          | 1807              | 0.32 |
| 16       | 32.94   | 222                | 41                | Ledol                       | 1924              | 0.42 |
| 17       | 34.95   | 220                | 41                | (-)-Caryophyllene- oxide    | 1977              | 1.62 |
| 18       | 35.70   | 222                | 41                | <i>cis</i> - Nerolidol      | 1996              | 0.90 |
| 19       | 36.61   | 222                | 43                | Epiglobulol                 | 2024              | 0.86 |
| 20       | 38.60   | 222                | 43                | Viridiflorol                | 2086              | 11   |
| 21       | 40.07   | 220                | 43                | Spathulenol                 | 2130              | 22.9 |
| 22       | 40.55   | 220                | 43                | Vulgarol A                  | 2144              | 1.65 |
| 23       | 41.70   | 222                | 43                | Globulol                    | 2176              | 1.16 |
| 24       | 43.23   | 220                | 43                | Aromadendren-epoxide-(I)    | 2221              | 3.84 |
| 25       | 45.51   | 222                | 41                | Carotol                     | 2288              | 1.16 |

$R_t^*$ , retention time in minutes; B P<sup>\*\*</sup>, base peak; KI<sup>\*\*\*</sup>, calculated Kovat's index.



**Table 2: Results of cytotoxic activity of the essential oil of *S. rowleyanus* Jacobs**

| Cell line | IC <sub>50</sub> (µg/ml ) | IC <sub>10</sub> (µg/ml ) |
|-----------|---------------------------|---------------------------|
| U251      | 5                         | 9.5                       |
| MCF8      | 2.6                       | 8                         |

*U251* = Brain tumor cell line.

*MCF8* = Liver carcinoma cell line.

IC<sub>50</sub> = Dose of the drug which reduces survival to 50%

IC<sub>10</sub> = Dose of the drug which reduces survival to 10%

**Table 3 : Results of the antimicrobial testing of the essential oil of *S. rowleyanus*.**

| Microorganism                           | Concentration % | Zone of inhibition | Rifampicin | Nystatin |
|-----------------------------------------|-----------------|--------------------|------------|----------|
| <i>Bacillus subtilis</i> ATCC 6633      | 40              | + + + +            |            |          |
|                                         | 20              | + + +              |            |          |
|                                         | 10              | + +                | + + +      | -        |
|                                         | 5               | -                  |            |          |
| <i>Staphylococcus aureus</i> ATCC 5141  | 40              | + + + +            |            |          |
|                                         | 20              | + + +              |            |          |
|                                         | 10              | + +                | + + + +    | -        |
|                                         | 5               | -                  |            |          |
| <i>Klebsiella pneumonia</i>             | 40              | + + +              |            |          |
|                                         | 20              | + +                |            |          |
|                                         | 10              | +                  | + + + +    | -        |
|                                         | 5               | -                  |            |          |
| <i>Escherichia coli</i> ATCC 10536      | 40              | + + +              |            |          |
|                                         | 20              | + +                |            |          |
|                                         | 10              | -                  | + + + +    | -        |
|                                         | 5               | -                  |            |          |
| <i>Proteus vulgaris</i> NCTC 4175       | 40              | + + +              |            |          |
|                                         | 20              | + +                |            |          |
|                                         | 10              | -                  | + + +      | -        |
|                                         | 5               | -                  |            |          |
| <i>Pseudomonas aeruginosa</i> NCTC 6750 | 40              | + + +              |            |          |
|                                         | 20              | -                  |            |          |
|                                         | 10              | -                  | + + +      | -        |
|                                         | 5               | -                  |            |          |
| <i>Candida albicans</i>                 | 40              | + + +              |            |          |
|                                         | 20              | + + +              |            |          |
|                                         | 10              | -                  | -          | + + +    |
|                                         | 5               | -                  |            |          |
| <i>Saccharomyces cerviciae</i>          | 40              | + + +              |            |          |
|                                         | 20              | + + +              |            |          |
|                                         | 10              | -                  | -          | + + +    |
|                                         | 5               | -                  |            |          |
| <i>Mycobacterium phlei</i> NRRL B-3683  | 40              | + + +              |            |          |
|                                         | 20              | + +                |            |          |
|                                         | 10              | +                  | -          | -        |
|                                         | 5               | -                  |            |          |

( - ) no zone of inhibition , + diameter of inhibition zone = 10 mm, + + diameter of inhibition zone = 11- 5 mm,  
 + + + diameter of inhibition zone = 11-22 mm, + + + + diameter of inhibition zone = 23-27 mm.

It is noteworthy to mention that to the best of our knowledge, this is the first report on the composition and bioactivities of the essential oil of *S. rowleyanus*.

#### REFERENCES

1. L H. Bailey and E. Z. Bailey. Hortus Third, p. 1170. Macmillan Publishing Co. New York, 1976
2. H. L. De Pooter, L. F. De Buyck, N. M. Schamp, E. A. Aboutabl, A. De Bruyn, and S. Z. Husain. The volatile fraction of *Senecio glaucus* sub sp. *Coronopifolius*. Flavor Fragrance Journal. **1**: 159 -163, 1986.
3. A.M. El Shazly. Essential oil composition of *Senecio desfontainei*. Zagazig J. Pharm. Sci. **8**: 1-8, 1999.
4. Z. G. Fernandes, V. I. Fernandez and G. B. Hammond Investigation of the essential oil of *Senecio tephrosioides* and *Salvia oppositiflora*. Revista- Latinoamericana-de Quimica. **25**: 14-16, 1996.
5. N., S. Mangi, N. Garg, S. K. Agarwal, and C. S. Mathela. The occurrence of  $\beta$ -thujone and a new p-menthane derivative in *Senecio chrysanthemoides* leaf oil. J. Ess. Oil Res. **7**: 511-514, 1995.
6. C. Perez, A. M. Agnese and J. L. Cabrera. The essential oil of *Senecio graveolens* (Compositae): Chemical composition and antimicrobial activity tests. J. Ethnopharmacol. **66**: 91-96, 1999.
7. B. van Dooren, R. Bos and D. H. E. Tattje. Composition of essential oils of some *Senecio* species. Planta Medica. **42**: 385-389, 1981.
8. R. R. Vera, S. J. Laurent and D. J. Fraisse. Chemical composition of the essential oil of *Senecio ambavilla* Bory Pers. from Reunion Island. J. Ess. Oil Res. **6**: 21, 1994.
9. M. A. Kuhnt, Pröbstle, G. Rimpler, R. Bauer, and M. Heinrich. Biological and pharmacological activities and further constituents of *Hyptis verticillata*. Planta Medica **61**: 227-232, 1995.
10. A. Sivoropouou, C. Nikolaou, E. Papanikolaou., S. Kokkini, T. Lanaras and M. Arsenakis. Antimicrobial, Cytotoxic and antiviral activities of *Salvia fruticosa* essential oil. J. Agric. Food Chem. **45**: 3197- 3201, 1997.
11. S.S. El-Hawary, Galal, A.M., Yousif, M.F. and Kirolos, F.N. Alkaloids and *in vitro* biological screening of *Senecio rowleyanus* Jacobs. Bull. Fac. Pharm. **44(3)**:185-192, 2007.
12. R. P. Adams. Identification of Essential Oil Components by Gas Chromatography / Mass Spectroscopy. Allured Publishing Corporation Carol Stream, Illinois, USA, 1995.
13. N. W. Davis, Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicon and Carbowax 20 M phases. J. Chromatography A. **503**: 1-24, 1990.
14. W. Jennings, and T. Shibamoto. Qualitative Analysis of Flavor and Fragrance Volatiles by Glass Capillary Gas Chromatography. Academic Press, London, 1980.
15. E. Kovats, Gas chromatographic characterization of organic substances in the retention index systems. Ch. 7: 229-247. In J. C. Giddings and R. A. Keller, **Advances in Chromatography**. Vol. 1. Marcel Dekker. Inc., New York, 1965.
16. P. Skehan, R. Storeng, D. Scudiero, A. Monks, J. McMahon, D. Vistica, J. T. Warren, H. Bokesch, S. Kenney, and M. R. Boyd. New colorimetric cytotoxicity assay for anticancer-drug screening. J Natl Cancer Inst. **82**: 1107-1112, 1990.
17. A. Monks, , D. Scudiero, P. Skehan, R. Shoemaker, K. Paull, D. Vistica, C. Hose, J. Langley, P. Cronise, A. Vaigro-Wolff, M. Gray-Goodrich, H. Campbell, J. Mayo, and M. Boyd. Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines. J. Natl Cancer Inst. **83**: 757-766, 1991.
18. J.P. Duguid, B.P. Marmion and R.H. A. Swain. **Medicinal Microbiology**, 13<sup>th</sup> ed., p.592. The English Language Book Society and Churchill, Livingstone London, 1978.
19. V. Lorian. Antibiotics in Laboratory Medicine 3rd ed, p. 134 – 144. Williams and Williams, Hong Kong, London, 1999.
20. M. Stephan, and J. A. Duke, **CRC Handbook of Medicinal Mints**. Library of Congress Inc., Boca Raton, New York, London and Tokyo, 1996.
21. M. E. El Tantawy. Chemical composition and biological activity of the essential oil of *Senecio micanioides* Otto cultivated in Egypt. Azh. J. Pharm. Sci., **26**:294-306, 2000.
22. M. R. Loizzo, R Statti, G.A., Tundis, R., Conforti, F., Bonesi, M., Autelitano, G., Houghton, P.J., Miljkovic-Brake, A., and Menichini, F.: Antibacterial and antifungal activity of *Senecio inaequidens* DC. and *Senecio vulgaris* L. Phytother. Res. **18(9)**:777-9, 2004. (Through <http://www.ncbi.nlm.nih.gov/pubmed/15478193>)
23. J. A. Duke, M. J. Bogenschutz-Godwin, J. duCellier, and P. K. Duke Handbook of Medicinal Herbs, 2<sup>nd</sup> Ed, CRC Press, Washington D.C., 2002.

Phcog Mag. Vol 4, Issue 16, Oct-Dec, 2008

Submitted on : 26<sup>th</sup> February, 2008

Revised on : 13<sup>th</sup> July, 2008

Accepted on : 17<sup>th</sup> July, 2008

\*\*\*\*\*